



# Hydrogen gas treatment prolongs replicative lifespan of bone marrow multipotential stromal cells *in vitro* while preserving differentiation and paracrine potentials

Haruhisa Kawasaki<sup>a,b</sup>, Jianjun Guan<sup>d</sup>, Kenichi Tamama<sup>a,b,c,\*</sup>

<sup>a</sup> Department of Pathology, The Ohio State University, Columbus, OH 43210, United States

<sup>b</sup> Davis Heart and Lung Research Institute, The Ohio State University, Columbus, OH 43210, United States

<sup>c</sup> Center for Stem Cell and Regenerative Medicine, The Ohio State University, Columbus, OH 43210, United States

<sup>d</sup> Department of Materials Science and Engineering, The Ohio State University, Columbus, OH 43210, United States

## ARTICLE INFO

### Article history:

Received 1 June 2010

Available online 4 June 2010

### Keywords:

Mesenchymal stem cells

Hydrogen gas

Senescence

Differentiation

Paracrine

Hypoxia

## ABSTRACT

Cell therapy with bone marrow multipotential stromal cells/mesenchymal stem cells (MSCs) represents a promising approach in the field of regenerative medicine. Low frequency of MSCs in adult bone marrow necessitates *ex vivo* expansion of MSCs after harvest; however, such a manipulation causes cellular senescence with loss of differentiation, proliferative, and therapeutic potentials of MSCs. Hydrogen molecules have been shown to exert organ protective effects through selective reduction of hydroxyl radicals. As oxidative stress is one of the key insults promoting cell senescence *in vivo* as well as *in vitro*, we hypothesized that hydrogen molecules prevent senescent process during MSC expansion. Addition of 3% hydrogen gas enhanced preservation of colony forming early progenitor cells within MSC preparation and prolonged the *in vitro* replicative lifespan of MSCs without losing differentiation potentials and paracrine capabilities. Interestingly, 3% hydrogen gas treatment did not decrease hydroxyl radical, protein carbonyl, and 8-hydroxydeoxyguanosine, suggesting that scavenging hydroxyl radical might not be responsible for these effects of hydrogen gas in this study.

© 2010 Elsevier Inc. All rights reserved.

## 1. Introduction

Adult bone marrow multipotential stromal cells/mesenchymal stem cells (MSCs) are multipotent cells with strong secretory activities of various growth factors. MSC-based cell therapy represents a promising approach to promote wound healing and tissue regeneration [1–9].

Sufficient number of MSCs retaining their multipotency and paracrine activity is needed for successful MSC-based therapeutics [10]. However, the frequency of MSCs declines age-dependently [11] so that these cells exist only one in  $10^5$ – $10^6$  adult bone marrow mononuclear cells [5]. *Ex vivo* MSC expansion could compensate for low harvest of MSCs; however, such an expansion causes cellular senescence with loss of differentiation, proliferative, and therapeutic potentials of MSCs [12–14].

Hydrogen molecules have been shown to exert organ protective effects in the ischemia–reperfusion injuries and post-transplant rejections through selective reduction of hydroxyl radicals and

possibly other unidentified mechanisms [15–18]. As oxidative stress is one of the key insults promoting cell senescence *in vivo* as well as *in vitro* [19], we hypothesized that hydrogen molecules prevent senescent process during *ex vivo* MSC expansion through anti-oxidative and cytoprotective effects. The aim of the present study was to test the hypothesis through addressing the effects of hydrogen molecules on replicative cell senescence, differentiation, and paracrine potentials of MSC.

## 2. Materials and methods

### 2.1. Cell culture

Cultured human primary bone marrow MSCs were from Lonza (Basel, Switzerland) for donor 1 and from Center for Gene Therapy, Tulane University (New Orleans, LA) for donor 2. Donor 1 was a 20 years old female and donor 2 was 19 years old male according to the product inserts. FBS was from Atlanta Biologicals (Lawrenceville, GA). MSCs were cultured in MEM $\alpha$  supplemented with 17% FBS, 2 mM L-glutamine, 1 mM pyruvate, and 100  $\mu$ M nonessential amino acids. Cell culture media and supplements were all from Invitrogen (Carlsbad, CA) unless otherwise stated. All of the

\* Corresponding author. Address: 320 West 10th Avenue, M364C Starling Loving Hall, Columbus, OH 43210, United States. Fax: +1 614 366 1183.

E-mail addresses: [kenichi.tamama@osumc.edu](mailto:kenichi.tamama@osumc.edu), [kjtamama@hotmail.com](mailto:kjtamama@hotmail.com) (K. Tamama).

experiments were conducted at 37 °C in a humidified air with 5% CO<sub>2</sub> unless otherwise specified.

## 2.2. Hydrogen gas treatment

Hydrogen gas treatment was made by culturing cells in premixed gas (3% H<sub>2</sub>, 21% O<sub>2</sub>, 5% CO<sub>2</sub>, balance N<sub>2</sub>) (Praxair, Danbury, CT) in a hypoxia chamber (Stemcell Technologies, Vancouver, Canada). In brief, cell culture dishes or multiwell plates were placed in the chamber equipped with airtight seal. Then, the chamber was flushed for 5 min or more with premixed gas (20 L/min) according to the manufacture's instruction. The chamber was flushed with premixed gas every 3 days or less to ensure the composition of mixed gas.

## 2.3. Hypoxia treatment

Hypoxic treatment was made by culturing cells in premixed gas (1% O<sub>2</sub>, 5% CO<sub>2</sub>, balance N<sub>2</sub>) (Praxair, Danbury, CT) in a hypoxia chamber (Stemcell Technologies, Vancouver, Canada) in a similar way to hydrogen gas treatment described previously.

## 2.4. Colony formation assay

Total of 500 cells was seeded in 100 mm culture dish and culture in 14 days as indicated. To exclude the possibility that hydrogen gas exposure affects MSC colony formation through altering adherence efficiency of MSC to plastic, MSCs were allowed to adhere to plastic after seeding for 24 h before initiation of hydrogen gas treatment. Upon completion of the indicated treatment, the formed colonies were rinsed with PBS twice to dislodge the dead cells and debris, then fixed and stained by 5% crystal violet in methanol. The number of colony formation (>2 mm diameter) was enumerated manually by three independent researchers. The results were given as a colony count per 1000 seeded cells.

## 2.5. Osteogenic differentiation

Osteogenic differentiation was induced as previously described [3]. In brief, MSCs were seeded at  $3 \times 10^4$  cells per 6-well and cultured in osteogenic differentiation medium containing dexamethasone, ascorbate, and  $\beta$ -glycerophosphate (Lonza, Basal, Switzerland) for 7 days. Osteogenic differentiation was evaluated by the induction of alkaline phosphatase expression, an early stage of osteogenic differentiation of MSC [20]. Intracellular alkaline phosphatase enzyme activity was detected by using quantitative colorimetric assay kit (BioAssay Systems, Hayward, CA) as well as by histochemical staining with StemTAG™ Alkaline Phosphatase Staining kit (Cell Biolab, San Diego, CA), according to manufactures' protocols.

## 2.6. Adipogenic differentiation

Adipogenic differentiation was induced as previously described after slight modifications [3]. In brief, MSCs were seeded at  $1 \times 10^6$  cells per 6-well and cultured in Adipogenic differentiation induction medium for 2–3 days followed by Adipogenic differentiation maintenance medium for 2 days in 1–2 cycles, as manufacture's instruction (Lonza, Basal, Switzerland). Adipogenic differentiation was evaluated by staining intracellular lipid droplets by Oil Red O and by induction of adipogenic marker peroxisome proliferation-activated receptor  $\gamma$  (PPAR $\gamma$ ) transcript [3,5].

## 2.7. Quantitative reverse transcription-polymerase chain reaction (qRT-PCR)

The level of mRNA transcripts was assessed by two-step qRT-PCR. In brief, after isolation of total RNA with TRIzol plus

(Invitrogen, Carlsbad, CA), cDNA was synthesized with High Capacity RNA-to-DNA kit (Applied Biosystems, Foster City, CA). Then, cDNA was mixed with Taqman® Universal Master Mix and each Taqman® gene-specific probe/primers (Applied Biosystems) and subject to the Taqman® PCR universal thermal cycling conditions defined by Applied Biosystems: 95 °C for 10 min and then, 40 cycles of two-temperature PCR at 95 °C for 15 s for denaturing, 60 °C for 1 min for annealing and extension. 7900MT (Applied Biosystems) was used as a quantitative real-time PCR thermal cycler.

Taqman® probe/primers were used for p16<sup>INK4A</sup> or cyclin-dependent kinase inhibitor 2A (Hs00923894\_m1), PPAR $\gamma$  (Hs01115513\_m1), indoleamine 2,3-dioxygenase-1 (IDO) (Hs00984148\_m1), and human ribosomal protein, large, P0 (RPLPO) (house-keeping gene) (4333761F) (Applied Biosystems). All PCR primers were designed to span intron(s) to discriminate cDNA amplicons from genomic amplicons. Comparative Ct method was utilized to assess the levels of each mRNA transcript relative to that level of RPLPO mRNA transcript.

## 2.8. Proliferation assay

MSCs were seeded in 6-well plate at the density of 50 cells/cm<sup>2</sup> (480 cells/well) in regular medium containing 17% FBS. Cell culture medium was changed every 3–4 days. Live cell counts were determined by manual enumeration of cells excluding trypan blue in the indicated timing, and the 480 live cells were seeded for the next round of experiment. The same process was repeated 3 times during 56 or 58-day experiments (Fig. 1C and D). Population doubling was calculated based upon those cell counts.

## 2.9. Quantitation of secreted growth factors, cytokine, and bioactive mediator


After rinsing with PBS, MSCs were cultured in 0.1% BSA-containing serum free medium for 24 h during the indicated treatments. The conditioned media were collected and the secreted VEGF, hepatocyte growth factor (HGF), basic fibroblast growth factor (bFGF), stromal cell-derived factor-1 (SDF-1), prostaglandin E<sub>2</sub> (PGE<sub>2</sub>), and interleukin-6 (IL-6) were measured by quantitative ELISA kits according to each manufactures' instructions (R&D Systems, Minneapolis, MN)(Cayman Chemical, Ann Arbor, MI). The readouts were adjusted per  $1 \times 10^6$  cells.

## 2.10. $\beta$ -Galactosidase staining

Senescence-associated  $\beta$ -galactosidase (SA- $\beta$ -gal) activity was detected histochemically by using Senescence  $\beta$ -Galactosidase Staining Kit according to the product insert (Cell Signaling Technology, Danvers, MA). SA- $\beta$ -gal positive cells were counted manually under microscope and expressed per 100 total cells.

## 2.11. Determination of intracellular reactive oxygen species (ROS)

The levels of intracellular ROS production were determined using 3-(p-hydroxyphenyl) fluorescein (HPF) (Invitrogen, Carlsbad, CA). Briefly, cell suspension ( $5 \times 10^5$  cells/ml) in PBS with 5% FBS was made after indicated treatment, and incubated with 5  $\mu$ M HPF for 30 min at 37 °C. All buffers and reagents were pre-equilibrated with the same mixed gas used in the treatment to prevent ROS production during preparation of cell suspension and incubation with HPF. The fluorochrome-loaded cells were excited using a 488 nm argon-ion laser and the fluorescent emission from HPF was recorded at 515–540 nm. Data were collected using BD FACSCalibur flowcytometry (BD Bioscience, Franklin Lakes, NJ) from at least 10,000 cells, and average signal strength was calculated per cell.



**Fig. 1.** (A and B) Effects of H<sub>2</sub> exposure on MSC colony formation. Five hundred cells were seeded in 10 cm dish and the number of formed colonies was counted in day 14 [ $p < 0.05$  to control (Ctrl) condition in (B)]. In (A), shown are representative images of MSC colony formation in the presence and absence of H<sub>2</sub>. (C and D) Effects of H<sub>2</sub> exposure on the replicative lifespan of MSC from donor 1 (C) and donor 2 (D) [ $**p < 1 \times 10^{-5}$ ,  $*p < 1 \times 10^{-3}$  to control (Ctrl) condition in (C) and (D), respectively]. (E–G) Effects of H<sub>2</sub> exposure on the induction of p16<sup>INK4A</sup> transcript by quantitative RT-PCR in (E) and SA-β-gal activity in (F) in donor 1 [ $*p < 0.005$  to control (Ctrl) in (E) and (F)]. In (G), shown are representative images of SA-β-gal staining of MSCs at day 58. Each photograph is 550 μm square.


## 2.12. Determination of oxidative damages

Oxidative DNA damage was assessed through quantitation of 8-hydroxydeoxyguanosine (8-OHdG) within cellular DNA by using

OxiSelect™ Oxidative DNA Damage ELISA kit (Cell Biolab). Protein carbonylation, the irreversible and unrepairable oxidative damage of protein [21], was evaluated through quantitation of protein carbonyls within protein lysates in Laemmli buffer by using OxiSelect™ Protein Carbonyl ELISA kit (Cell Biolab).

## 2.13. Statistical analysis

All experiments were performed in duplicate or triplicate. Data were analyzed using student *t*-tests. Significance was set at  $P < 0.05$  or more stringent as noted in the text and figure legends. The data of multiple observations were provided as mean  $\pm$  SEM for at least three separate experiments unless stated otherwise.



**Fig. 2.** Effects of H<sub>2</sub> exposure on the osteogenic and adipogenic differentiation of MSCs. (A, C, and E) Effects of H<sub>2</sub> exposure on osteogenic differentiation of MSCs. Cells were cultured in regular medium (Reg) or osteogenic medium (Osteo) in the presence and absence of H<sub>2</sub> for 7 days, and an early phase of osteogenic differentiation was evaluated by histochemical staining of alkaline phosphatase macroscopically in (A) and microscopically in (C) and biochemical quantitation of alkaline phosphatase activity in (E) [ $*p < 1 \times 10^{-6}$ ,  $#p < 1 \times 10^{-6}$  to regular medium control of each respective condition in (E)]. (B, D, and F) Effects of H<sub>2</sub> exposure on adipogenic differentiation of MSCs. Cells were cultured in regular medium (Reg) or 2 cycles of adipogenic stimulations (Adipo) in the presence and absence of H<sub>2</sub>, and an adipogenic differentiation was evaluated with Oil Red O staining of intracellular lipid droplets macroscopically (B) and microscopically (D) and with mRNA transcript levels of PPARγ by quantitative RT-PCR (F) [ $*p < 0.005$ ,  $#p < 0.01$  to regular medium control of each respective condition in (F)]. Each photograph is 550 μm square in (E) and (F).


### 3. Results

#### 3.1. Hydrogen gas treatment preserves colony forming early progenitors within MSCs population and prolongs replicative lifespan of MSCs *in vitro*

One of the prominent characteristics of MSC is their ability to produce colonies after seeded at low density [6,22]. MSC populations are shown to be heterogeneous *in vitro*, containing both colony forming

early progenitors or rapidly self-renewing (RS) cells and mature/senescent cells [12,23,24]. Preservation of the colony forming early progenitors is critical during *ex vivo* MSC expansion for successful MSC-based therapeutics, which can be assessed easily by colony forming assay [12,22,25]. Hydrogen gas treatment increased colony formation of MSCs, suggesting that early progenitor fraction of MSCs is better preserved with hydrogen gas treatment (Fig. 1A and B).

Prolongation of replicative lifespan is another important mechanism to promote *ex vivo* MSC expansion. Oxidative stress is



**Fig. 3.** Effects of H<sub>2</sub> exposure on the secretion of bFGF (A), HGF (B), VEGF (C), SDF-1 (D), IL-6 (E), PGE<sub>2</sub> (F and G), and IDO (H and I) from MSCs. For H<sub>2</sub>(+) group, cells were incubated with H<sub>2</sub> for 4 days before collection of conditioned media. For H<sub>2</sub> pretreatment (preTx) group, cells were incubated with H<sub>2</sub> for 4 days, and then cultured for additional 4 days without H<sub>2</sub> before collection of conditioned media. The levels of growth factors and cytokines in the serum-starved conditioned media were determined by ELISA. The levels of IDO mRNA transcript relative to RPLPO transcripts (housekeeping gene) were by quantitative RT-PCR (H and I). The levels of PGE<sub>2</sub> and IDO were determined in the presence (G and I) and absence (F and H) of 20 ng/mL interferon-γ (INF-γ). *p*-Values were given in each figure.



regarded one of the main mechanisms limiting replicative lifespan through accelerating senescent process [19]. Hydrogen gas treatment prolonged replicative lifespan of culture MSCs obtained from two independent donors (Fig. 1C and D).

SA- $\beta$ -gal is a well-known marker widely used to identify senescent cells [26]. p16<sup>INK4A</sup> expression has been shown to elevate in senescent MSCs [27]. Hydrogen gas delayed cellular senescence process, as it lowered the number of MSCs expressing SA- $\beta$ -gal and the overall transcript level of p16<sup>INK4A</sup> at day 58 of culture (Fig. 1E–G).

### 3.2. Hydrogen gas treatment preserves differentiation potentials of MSCs

Multidifferentiation potential of MSC is another indispensable feature for MSC-based therapeutics and tissue engineering, and it should be preserved during *ex vivo* MSC expansion. Osteogenic and adipogenic differentiations are two major differentiation fates for MSC [3–5]. Hydrogen gas treatment did not induce either osteogenic or adipogenic differentiations, and it did not alter both differentiations induced by osteogenic and adipogenic differentiation media, respectively (Fig. 2).

### 3.3. Hydrogen gas treatment preserves paracrine potentials of MSCs

MSC transplantation was shown to promote angiogenesis and tissue regeneration through its strong paracrine capability of various growth factors such as VEGF, bFGF, or HGF [7,28]. MSCs were also shown to exert anti-inflammatory or immunomodulatory effects through paracrine mechanisms [29,30]. Thus, it is critically important to evaluate the effects of hydrogen gas treatment on MSC paracrine profile.

Hydrogen gas treatment increased the secretion of bFGF and HGF, whereas it decreased VEGF secretion (Fig. 3A–C). SDF-1 is a chemokine which attracts stem/progenitor cells to regenerating tissues [31,32], and SDF-1-mediated recruitment of stem/progenitor cells is believed to be one of the mechanisms in MSC-mediated angiogenesis [33]. SDF-1 secretion from MSCs was unaltered by hydrogen gas treatment (Fig. 3D). IL-6 is a pro-inflammatory cytokine, but at the same time, it was also shown to play a key role in self-renewal of undifferentiated MSCs *in vitro* [34]. IL-6 secretion was decreased by hydrogen gas treatment (Fig. 3E).

PGE<sub>2</sub> and IDO are the ones of the main molecules mediating anti-immunologic and anti-inflammatory effects of MSCs [30]. Interferon- $\gamma$  (IFN- $\gamma$ ) was shown to enhance immunosuppressive

property of MSC through up-regulation of IDO [35,36]. Although hydrogen gas treatment decreased the PGE<sub>2</sub> secretion, it increased IDO expression in MSCs; however, such an effect was not apparent in the presence of IFN- $\gamma$ , presumably due to extremely strong induction of IDO by IFN- $\gamma$  (Fig. 3F–I).

### 3.4. Hydrogen gas might exert these effects without scavenging hydroxyl radicals


Hydrogen molecule was initially shown to exert anti-oxidant effect through selectively reducing hydroxyl radical, the most powerful ROS in the body [18]; however, the involvements of other unknown mechanisms have been suggested for hydrogen-associated cytoprotective effects [37].

We chose 3% as a concentration of hydrogen gas because of serious and realistic concerns about flammability of hydrogen gas over 4.6%; however, 3% is much lower than the one used in the initial study describing hydroxyl radical scavenging effects by hydrogen molecule [18]. Interestingly, 3% hydrogen gas did not reduce the fluorescent intensity of HPF, whereas 1% O<sub>2</sub> decreased it (Fig. 4A). Moreover, it did not decrease the levels of protein carbonyls and 8-OHdG, whereas 1% O<sub>2</sub> decreased both of them (Fig. 4B and C). These data suggested that anti-oxidant effect might not be responsible for these hydrogen gas-mediated effects in this study.

## 4. Discussion

Oxidative stress is one of the major insults accelerating cell senescence *in vivo* as well as *in vitro* [19]. Reduction of oxidative stress by lowering oxygen tension or adding anti-oxidant such as vitamin C or N-acetylcysteine has been shown to prolong replicative lifespan of human cells including MSCs *in vitro* [38–41]. Hydrogen molecule was shown to exert anti-oxidant effect through selectively reducing hydroxyl radical, the most powerful ROS in the body [18], and indeed, hydrogen gas prolonged the *in vitro* replicative lifespan of MSCs without losing differentiation potentials and paracrine capabilities. Hydrogen gas treatment also enhanced preservation of colony forming early progenitor cells within MSC preparation.

However, 3% hydrogen gas treatment did not decrease hydroxyl radical, protein oxidative carbonylation and 8-OHdG (Fig. 4). The apparent mechanism is unclear at this moment, but it indicates that 3% hydrogen gas might be too low to scavenge hydroxyl radical, but still adequate to enhance MSC expansion *ex vivo*. Indeed,



**Fig. 4.** Effects of H<sub>2</sub> or hypoxia (1% O<sub>2</sub>) on the generation of intracellular reactive oxygen species (ROS) (A), 8-OHdG (B), and protein carbonyls (C). MSCs were cultured in the presence of H<sub>2</sub> or under hypoxic condition for 10 days prior to assays. Intracellular ROS levels were determined using fluorescent intensity of HPF by flowcytometry. The levels of 8-OHdG and protein carbonyls were determined by ELISA [ $p < 0.005$  (A),  $*p < 0.05$  (B and C) to 20% O<sub>2</sub> condition in each graph].

the involvement of the unknown mechanisms has previously been suggested [37], which is consistent with our data.

Hydrogen gas treatment did not diminish paracrine activity of MSCs; still, it altered the paracrine profiles (Fig. 3). For example, hydrogen gas treatment increased both bFGF and HGF secretions whereas it decreased VEGF secretion. Hydrogen gas treatment increased the IDO expression, whereas it decreased PGE<sub>2</sub>. The net effects of hydrogen gas treatment on the MSC-mediated angiogenic and anti-inflammatory/immunomodulatory actions are uncertain based on these data alone. The effects of hydrogen gas treatment appeared reversible, though some residual effects of hydrogen gas was still observed 5 days after it was discontinued (Fig. 3), suggesting that pretreatment of MSCs with hydrogen gas might still be effective after MSC transplantation *in vivo*.

In conclusion, we demonstrate that hydrogen gas treatment enhances preservation of colony forming early progenitor cells within MSC preparation and prolongs the *in vitro* replicative lifespan of MSCs without losing differentiation potentials and paracrine capabilities. Hydroxyl radical scavenging effects might not be responsible for these effects of 3% hydrogen gas.

## Acknowledgments

This study was supported by AHA Beginner-Grant-in-aid (09BGIA2050227) (K.T.). Some of the materials employed in this work were provided by the Tulane Center for Gene Therapy through a grant from NIH (P40RR017447). We would like to thank Dr. Darwin J. Prockop (Tulane Center for Gene Therapy, New Orleans, LA, currently at Texas A&M Health Science Center, Temple, TX) for providing the primary human MSCs.

## References

- [1] K. Tamama, H. Kawasaki, A. Wells, Epidermal growth factor (EGF) treatment on multipotential stromal cells (MSCs), possible enhancement of therapeutic potential of MSC, J. Biomed. Biotechnol. 2010 (2010) 795385.
- [2] K. Tamama, C.K. Sen, A. Wells, Differentiation of bone marrow mesenchymal stem cells into the smooth muscle lineage by blocking ERK/MAPK signaling pathway, Stem Cells Dev. 17 (2008) 897–908.
- [3] K. Tamama, V.H. Fan, L.G. Griffith, H.C. Blair, A. Wells, Epidermal growth factor as a candidate for ex vivo expansion of bone marrow-derived mesenchymal stem cells, Stem cells 24 (2006) 686–695.
- [4] D.J. Prockop, Marrow stromal cells as stem cells for nonhematopoietic tissues, Science 276 (1997) 71–74.
- [5] M.F. Pittenger, A.M. Mackay, S.C. Beck, R.K. Jaiswal, R. Douglas, J.D. Mosca, M.A. Moorman, D.W. Simonetti, S. Craig, D.R. Marshak, Multilineage potential of adult human mesenchymal stem cells, Science 284 (1999) 143–147.
- [6] A.J. Friedenstein, J.F. Gorskaja, N.N. Kulagina, Fibroblast precursors in normal and irradiated mouse hematopoietic organs, Exp. Hematol. 4 (1976) 267–274.
- [7] T. Kinnaird, E. Stabile, M.S. Burnett, S.E. Epstein, Bone marrow-derived cells for enhancing collateral development: mechanisms, animal data, and initial clinical experiences, Circ. Res. 95 (2004) 354–363.
- [8] M.F. Pittenger, B.J. Martin, Mesenchymal stem cells and their potential as cardiac therapeutics, Circ. Res. 95 (2004) 9–20.
- [9] Y. Wu, L. Chen, P.G. Scott, E.E. Tredget, Mesenchymal stem cells enhance wound healing through differentiation and angiogenesis, Stem cells 25 (2007) 2648–2659.
- [10] V. Falanga, S. Iwamoto, M. Chartier, T. Yufit, J. Butmarc, N. Kouttab, D. Shrayar, P. Carson, Autologous bone marrow-derived cultured mesenchymal stem cells delivered in a fibrin spray accelerate healing in murine and human cutaneous wounds, Tissue Eng. 13 (2007) 1299–1312.
- [11] A.I. Caplan, Why are MSCs therapeutic? New data: new insight, J. Pathol. 217 (2009) 318–324.
- [12] D.C. Colter, R. Class, C.M. DiGirolamo, D.J. Prockop, Rapid expansion of recycling stem cells in cultures of plastic-adherent cells from human bone marrow, Proc. Natl. Acad. Sci. USA 97 (2000) 3213–3218.
- [13] P.R. Crisostomo, M. Wang, G.M. Wairiuko, E.D. Morrell, A.M. Terrell, P. Seshadri, U.H. Nam, D.R. Meldrum, High passage number of stem cells adversely affects stem cell activation and myocardial protection, Shock 26 (2006) 575–580.
- [14] V.D. Roobrouck, F. Ulloa-Montoya, C.M. Verfaillie, Self-renewal and differentiation capacity of young and aged stem cells, Exp. Cell Res. 314 (2008) 1937–1944.
- [15] B.M. Buchholz, D.J. Kaczorowski, R. Sugimoto, R. Yang, Y. Wang, T.R. Billiar, K.R. McCurry, A.J. Bauer, A. Nakao, Hydrogen inhalation ameliorates oxidative stress in transplantation induced intestinal graft injury, Am. J. Transplant. 8 (2008) 2015–2024.
- [16] J.S. Cardinal, J. Zhan, Y. Wang, R. Sugimoto, A. Tsung, K.R. McCurry, T.R. Billiar, A. Nakao, Oral hydrogen water prevents chronic allograft nephropathy in rats, Kidney Int. 77 (2010) 101–109.
- [17] K. Hayashida, M. Sano, I. Ohsawa, K. Shinmura, K. Tamaki, K. Kimura, J. Endo, T. Katayama, A. Kawamura, S. Kohsaka, S. Makino, S. Ohta, S. Ogawa, K. Fukuda, Inhalation of hydrogen gas reduces infarct size in the rat model of myocardial ischemia-reperfusion injury, Biochem. Biophys. Res. Commun. 373 (2008) 30–35.
- [18] I. Ohsawa, M. Ishikawa, K. Takahashi, M. Watanabe, K. Nishimaki, K. Yamagata, K. Katsura, Y. Katayama, S. Asoh, S. Ohta, Hydrogen acts as a therapeutic antioxidant by selectively reducing cytotoxic oxygen radicals, Nat. Med. 13 (2007) 688–694.
- [19] J. Campisi, From cells to organisms: can we learn about aging from cells in culture?, Exp. Gerontol. 36 (2001) 607–618.
- [20] N. Jaiswal, S.E. Haynesworth, A.I. Caplan, S.P. Bruder, Osteogenic differentiation of purified, culture-expanded human mesenchymal stem cells in vitro, J. Cell. Biochem. 64 (1997) 295–312.
- [21] T. Nystrom, Role of oxidative carbonylation in protein quality control and senescence, EMBO J. 24 (2005) 1311–1317.
- [22] R. Pochampally, Colony forming unit assays for MSCs, Methods Mol. Biol. 449 (2008) 83–91.
- [23] D.C. Colter, I. Sekiya, D.J. Prockop, Identification of a subpopulation of rapidly self-renewing and multipotential adult stem cells in colonies of human marrow stromal cells, Proc. Natl. Acad. Sci. USA 98 (2001) 7841–7845.
- [24] I. Sekiya, B.L. Larson, J.R. Smith, R. Pochampally, J.G. Cui, D.J. Prockop, Expansion of human adult stem cells from bone marrow stroma: conditions that maximize the yields of early progenitors and evaluate their quality, Stem Cells 20 (2002) 530–541.
- [25] J.R. Smith, R. Pochampally, A. Perry, S.C. Hsu, D.J. Prockop, Isolation of a highly clonogenic and multipotential subfraction of adult stem cells from bone marrow stroma, Stem Cells 22 (2004) 823–831.
- [26] G.P. Dimri, X. Lee, G. Basile, M. Acosta, G. Scott, C. Roskelley, E.E. Medrano, M. Linskens, I. Rubelj, O. Pereira-Smith, et al., A biomarker that identifies senescent human cells in culture and in aging skin in vivo, Proc. Natl. Acad. Sci. USA 92 (1995) 9363–9367.
- [27] K.R. Shibata, T. Aoyama, Y. Shima, K. Fukiage, S. Otsuka, M. Furu, Y. Kohno, K. Ito, S. Fujibayashi, M. Neo, T. Nakayama, T. Nakamura, J. Toguchida, Expression of the p16INK4A gene is associated closely with senescence of human mesenchymal stem cells and is potentially silenced by DNA methylation during in vitro expansion, Stem Cells 25 (2007) 2371–2382.
- [28] L. Chen, E.E. Tredget, P.Y. Wu, Y. Wu, Paracrine factors of mesenchymal stem cells recruit macrophages and endothelial lineage cells and enhance wound healing, PLoS One 3 (2008) e1886.
- [29] S.S. Iyer, M. Rojas, Anti-inflammatory effects of mesenchymal stem cells: novel concept for future therapies, Expert Opin. Biol. Ther. 8 (2008) 569–581.
- [30] A. Uccelli, L. Moretta, V. Pistoia, Mesenchymal stem cells in health and disease, Nat. Rev. Immunol. 8 (2008) 726–736.
- [31] I. Petit, D. Jin, S. Rafii, The SDF-1-CXCR4 signaling pathway: a molecular hub modulating neo-angiogenesis, Trends Immunol. 28 (2007) 299–307.
- [32] K. Stellos, M. Gawaz, Platelets and stromal cell-derived factor-1 in progenitor cell recruitment, Semin. Thromb. Hemost. 33 (2007) 159–164.
- [33] H. Haider, S. Jiang, N.M. Idris, M. Ashraf, IGF-1-overexpressing mesenchymal stem cells accelerate bone marrow stem cell mobilization via paracrine activation of SDF-1alpha/CXCR4 signaling to promote myocardial repair, Circ. Res. 103 (2008) 1300–1308.
- [34] K.L. Pricola, N.Z. Kuhn, H. Haleem-Smith, Y. Song, R.S. Tuan, Interleukin-6 maintains bone marrow-derived mesenchymal stem cell stemness by an ERK1/2-dependent mechanism, J. Cell. Biochem. 108 (2009) 577–588.
- [35] M. Kramer, L. Cosmi, R. Angeli, A. Pasini, F. Liotta, A. Andreini, V. Santarlasci, B. Mazzinghi, G. Pizzolo, F. Vinante, P. Romagnani, E. Maggi, S. Romagnani, F. Annunziato, Role for interferon-gamma in the immunomodulatory activity of human bone marrow mesenchymal stem cells, Stem Cells 24 (2006) 386–398.
- [36] J.M. Ryan, F. Barry, J.M. Murphy, B.P. Mahon, Interferon-gamma does not break, but promotes the immunosuppressive capacity of adult human mesenchymal stem cells, Clin. Exp. Immunol. 149 (2007) 353–363.
- [37] K.C. Wood, M.T. Gladwin, The hydrogen highway to reperfusion therapy, Nat. Med. 13 (2007) 673–674.
- [38] G. Kashino, S. Kodama, Y. Nakayama, K. Suzuki, K. Fukase, M. Goto, M. Watanabe, Relief of oxidative stress by ascorbic acid delays cellular senescence of normal human and Werner syndrome fibroblast cells, Free Radic. Biol. Med. 35 (2003) 438–443.
- [39] C. Fehrer, R. Brunauer, G. Laschober, H. Unterluggauer, S. Reitingner, F. Kloss, C. Gully, R. Gassner, G. Lepperdinger, Reduced oxygen tension attenuates differentiation capacity of human mesenchymal stem cells and prolongs their lifespan, Aging Cell 6 (2007) 745–757.
- [40] T.M. Lin, J.L. Tsai, S.D. Lin, C.S. Lai, C.C. Chang, Accelerated growth and prolonged lifespan of adipose tissue-derived human mesenchymal stem cells in a medium using reduced calcium and antioxidants, Stem Cells Dev. 14 (2005) 92–102.
- [41] Y. Jin, T. Kato, M. Furu, A. Nasu, Y. Kajita, H. Mitsui, M. Ueda, T. Aoyama, T. Nakayama, T. Nakamura, J. Toguchida, Mesenchymal stem cells cultured under hypoxia escape from senescence via down-regulation of p16 and extracellular signal regulated kinase, Biochem. Biophys. Res. Commun. 391 (2010) 1471–1476.